

AD_____

Award Number: W81XWH-06-1-0650

TITLE: Seladin-1: A Novel Tumor Suppressor Gene Involved in Breast Cancer?

PRINCIPAL INVESTIGATOR: Konstantin I. Galaktionov, Ph.D.

CONTRACTING ORGANIZATION: Baylor College of Medicine
Houston, TX 77030

REPORT DATE: August 2008

TYPE OF REPORT: Final

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE 16-08-2008		2. REPORT TYPE Final		3. DATES COVERED 17 JUL 2006 - 16 JUL 2008	
4. TITLE AND SUBTITLE Seladin-1: A Novel Tumor Suppressor Gene Involved in Breast Cancer?				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER W81XWH-06-1-0650	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Konstantin I. Galaktionov, Ph.D. Email: kgalakt@bcm.tmc.edu				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Baylor College of Medicine Houston, TX 77030				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT Seladin-1 maps to the human chromosome region 1p31-1p32 that shows frequent loss of heterozygosity (LOH) in human breast tumors. Publicly available data (Entrez, GEO) also show a significant variability between levels of Seladin-1 expression in breast cancer cell lines and normal breast epithelium. Is Seladin-1 a tumor suppressor on 1p31-1p32 that is involved in breast cancer? Our concept is that it is. In order to determine if Seladin-1 is a TSG involved in breast cancer we propose to: 1) Identify possible Seladin-1 mutations in primary breast tumors. 2) Investigate the alterations of Seladin-1 expression in breast cancer cells. 3) Perform functional assays on tumor-specific Seladin-1 mutants. Accordingly, we amplified Seladin-1 exons from from 60 breast tumor genomic DNA samples (obtained from tissue banks or commercial sources), followed by the sequence analysis of the open reading frames. As a result, no missense or nonsense mutations were detected. We found that expression of Seladin-1 significantly varies between different breast cancer cell lines, and in one such line, MDA-MB-231, is significantly below expression in normal breast epithelium.					
15. SUBJECT TERMS breast cancer, breast cancer cell lines, Seladin-1					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT	b. ABSTRACT	c. THIS PAGE			USAMRMC
U	U	U	UU	10	19b. TELEPHONE NUMBER (include area code)

Table of Contents

	Page
Introduction.....	4
Body.....	4-7
Key Research Accomplishments.....	7
Reportable Outcomes.....	7
Conclusions.....	7
References.....	7-8
Appendices.....	9-10

INTRODUCTION:

Seladin-1 has been described recently as a novel regulator of p53, a major human tumor suppressor gene (TSG; [1]). Seladin-1 acts to counteract Mdm2 and promote accumulation of p53 in response to oncogenes and oxidative stress [1]. P53 is implicated in breast cancer progression, but it is mutated only in a fraction of human breast tumors [2], suggesting that additional mechanisms might be involved in functional inactivation of the p53 checkpoint in tumors expressing wild type p53. Additionally, Seladin-1 (under DHCR24 alias) is also involved in cholesterol metabolism [3]. Cholesterol metabolism is known to be a contributing factor in cancer progression.

In order to determine if Seladin-1 is a TSG involved in breast cancer we propose to: 1) Identify possible Seladin-1 mutations in primary breast tumors. 2) Investigate the alterations of Seladin-1 expression in breast cancer cells. 3) Perform functional assays on tumor-specific Seladin-1 mutants.

BODY:

Task1. Identify possible Seladin-1 mutations in primary breast tumors (months 1-12).

Initially, we amplified the Seladin-1 exons from 60 breast tumor genomic DNA samples (all were obtained from a commercial source (Oncomatrix, Inc.), followed by the sequence analysis of the open reading frames. As a result, we did not encounter finding mutations in the coding region and exon-intron junctions (other than known single nucleotide polymorphisms (SNPs).

However, we found high frequency (12-25%) new insertion/deletion polymorphism in the area close to exon7 (Fig.1). We observed a single nucleotide deletion (IVS6: delT -29...-36) in a stretch of seven consequent T upstream of exon 7 of Seladin-1 (see Fig.1) in genomic DNA of patients with breast tumors. This type of sequence belongs to a structure commonly referred to as the polypyrimidine tract. The exact size of polypyrimidine tract is gene-specific, but it has been shown that a single nucleotide deletion in this tract results in exon skipping or abnormal splicing, sometimes causing nonsense-mediated mRNA decay and resulting in several different inherited diseases.

Figure 1. In Pink: forward and reverse primer (remember that actual sequence is the reverse complement of what is shown for reverse primer only). Blue underlined is exon 7. Italics: a potential lariat branchpoint sequence. Stretch of T's with one nucleotide deleted (IVS6: delT –29...–36) in mutant is underlined.

Fr7fwd2 ➔ In mutant: TTTTTT

GCTTCTGTCAAGCACAGAATGGGACCTGGCCCCCTGGCCCCTTGCACTTGGAGACATTCAACCCTTGTTGGTTTTTTTTT

Exon7

CTCTCCAACCCCTAAATCTTTGTTTTCAGGACATTATCCCCCTTGGCAACAACCCCATCTTCCGCTACCTCTTTGGCTGGATGGTGCCTC
 CCAAGATCTCCCTCCTGAAGCTGACCCAGGGTGAGACCCTGCGCAAGCTGTACGAGCAGCACCAC
 GTGGTGCAAGACATGCTGGTGCCCATGAAGTGCCTGCAGCAGGCCCTGCACACCTTCCAAAACGACATCCACGTGAG


Fr7rev2 ←

TGGGCGAGGGCAGGGAGGGCGCACAGGTGCATTCCCTTGAATCCATGGTCTCCTCCCCTGAACCGTGTGACCGACCTTGGGCACACT
 GCCCTCTGGGTGTTTTCCCAACTTTAAGATGAGGTCCTTGTCCTATGTCTGCTGAGGCCTCAAC

The actual sequence analysis of one breast tumor sample with this polymorphism is shown in Fig. 2.

Figure 2. Mutation in human Seladin-1 intron-exon junction sequence (IVS6: delT –29...–36). Top-genomic DNA from primary human tumor (Reverse sequence shown), bottom-normal human DNA. Position of one nucleotide deletion (delT) is shown. Note that reverse complementary sequence is shown as indicated.

Reverse sequence



We analyzed whether the matching DNA from unaffected organs of the same patient. In case of this particular polymorphism, both tumor material and patients DNA obtained from unaffected tissues have the same polymorphism. We are now investigated whether this particular polymorphism is present with higher or lower frequency in breast cancer patient population comparing with unaffected individuals. No statistically significant differences were shown, therefore this polymorphism is unlikely to be important for the eventual development of breast cancer.

We submitted an abstract describing early stages of this research to Cold Spring Harbor Symposium on Quantitative Biology (see attached).

Our final analysis of this polymorphism was accepted to the Era of Hope Breast Cancer Conference (see attached).

Task2. Investigate alterations of Seladin-1 expression in breast cancer cells (months 1-12).

Accordingly, we isolated RNAs from human breast cell lines MCF7, BTK20, T47D, HBL100, MDA-MB-231, MDA-MB-468 and MCF10A. Total RNA isolation was done with Trizol (Invitrogen) as described by the manufacturer. Seladin-1 Probe labeling and hybridization was done as described previously using commercial kits. As a control, we used RNA from three different batches of normal mammary epithelial cells (HMEC). For immunoblotting, we used antibodies against C-terminal domain of Seladin-1, described in [1].

When RNAs from 10 primary breast cancer cell lines were analyzed, the conclusions on Seladin-1 gene expression was that this gene is usually expressed at high level in breast cancer cells (with the exception of MDA-MB-231). Interestingly, expression of Seladin-1 in MDA-MB-231 cell line was subjected to regulation by 5aza-2'-deoxycytidine (5AzadC), a demethylating agent. An increase in Seladin-1 expression in response to 5AzaDC indicates that this putative cell line was a candidate for promoter methylation studies using methylation-specific PCR and bisulfite sequencing. However, despite our best efforts, none of the primers designed to perform methylation studies had worked on promoter region of Seladin-1 (probably due to very high C/G content in those areas). More efforts in this regard are currently ongoing.

Task3. Perform functional assays on tumor-specific Seladin-1 mutants (months 7-12).

So far, we did not detect breast tumor-specific Seladin-1 mutants (in the coding region) and, therefore, we concentrated on mutations that our group [1] and others [3] have previously described. Interestingly, we found that Seladin-1 had inhibitory effect on growth of some breast cancer cells (notably MDA-MB-231) expressing it at the low levels, while having no affect on the others (T47D or MDA-MB-468) that express endogenous Seladin-1 at higher levels.

KEY RESEARCH ACCOMPLISHMENTS:

1. From analysis of 60 genomic DNA samples from patients with breast cancers, we did not find mutations in coding (exon) sequences, suggesting that a potential Seladin-1 mutation in breast cancer is a rare event.
2. We identified novel insertion/deletion polymorphism in a close proximity to exon 7 in Seladin-1 genomic sequence. We plan to analyze frequency of this polymorphism with relation to breast cancer and other malignancies.
3. Some breast cancer cell lines growth (MDA-MB-231) is inhibited by Seladin-1 in vitro.

REPORTABLE OUTCOMES:

1. ABSTRACT presented at the 70th Cold Spring Harbor Symposium on “Molecular approaches to controlling cancer”, 2005.
2. NIH RO1-A1 grant “Seladin-1 in RAS induced senescence and tumorigenesis” applied for in March of 2006.
3. Abstract accepted at Era of Hope Conference, Baltimore 2008.

CONCLUSIONS:

1. Seladin-1 is rarely mutated in breast cancer.
2. A novel high frequency (12-25%) Seladin-1 insertion/deletion polymorphism was found.
3. Seladin-1 inhibits growth of some breast cancer cell lines in vitro.

REFERENCES:

1. Wu, C., Miloslavskaya, I, Demontis, S., Maestro, R., and Galaktionov, K.. *Nature* 2004, **432**, 640-645.

2. Yamashita, H., Nishio, M., Toiyama, T., Sugiura, H., Zhang, Z, Kobayashi, S., and Iwase, H. Coexistence of HER2 over-expression and p53 protein accumulation is a strong prognostic molecular marker in breast cancer. *Breast Cancer Res.* 2004, **6**, R24-R30. Epub 2003 Nov 07.
3. Waterham, H.R. et al. Mutations in the 3beta-hydroxysterol Delta24-reductase gene cause desmosterolosis, an autosomal recessive disorder of cholesterol biosynthesis. *Am. J Hum. Genet.* 2001, **69**, 685-694.

SELADIN-1 MUTATION ASSOCIATED WITH MULTIPLE HUMAN TUMORS

Konstantin Galaktionov¹, Roberta Maestro², Olga Chernova³, Gene H. Barnett³, Madhuri Hegde¹, Silvia Demontis², and Irene Miloslavskaya¹.

¹ Department of Molecular and Human Genetics, Baylor College of Medicine, Houston, TX 77030 USA. ² MMNP Unit-Department of Experimental Oncology CRO-IRCCS, Aviano National Cancer Institute, 33081 Aviano (PN), Italy. ³ Brain Tumor Institute, The Cleveland Clinic Foundation, Cleveland, USA

Expression of multiple oncogenes and inactivation of tumor suppressors is required to transform primary mammalian cells into cancer cells. Activated Ha-*RasV12* (Ras) is usually associated with cancer, but it also produces paradoxical premature senescence in primary cells by inducing reactive oxygen species followed by accumulation of tumor suppressors p53 and p16^{INK4a}. We identified, using a direct genetic screen, *Seladin-1* as a key mediator of Ras-induced senescence (Wu *et al.* 2004, *Nature* 432, 640). Following oncogenic and oxidative stress, Seladin-1 binds p53 amino terminus and displaces E3 ubiquitin ligase Mdm2 from p53, thus resulting in p53 accumulation. This suggested that *Seladin-1* is a potential human tumor suppressor.

In order to test this hypothesis, we performed the sequence analysis of coding exons of Seladin-1 in 97 human sporadic cancers of various types. As the result, we observed that approximately 21% of all tested patients (20 out of 97) had the mutation at the single common site, suggesting that *Seladin-1* function is frequently affected in patients with multiple major tumor types. No known SNPs exist at this location and those were not detected at this site in 64 randomly chosen unaffected donors. The cancer mutation data and the significance of this mutation for Seladin-1 function will be presented.

The remarkably high frequency of the *Seladin-1* single-site mutation in patients developing multiple human tumors suggests a potentially unique requirement for, or a mechanism of, *Seladin-1* mutation in human cancers in comparison with the previously described tumor suppressors.

Seladin-1 has been described as a novel regulator of p53, a major human tumor suppressor gene (TSG; [1]). Seladin-1 acts to counteract Mdm2 and promote accumulation of p53 in response to oncogenes and oxidative stress [1]. P53 is implicated in breast cancer progression, but it is mutated only in a fraction of human breast tumors [2], suggesting that additional mechanisms might be involved in functional inactivation of the p53 checkpoint in tumors expressing wild type p53. In order to determine if Seladin-1 is a TSG involved in breast cancer we proposed to: 1) Identify possible Seladin-1 mutations in primary breast tumors. 2) Investigate the alterations of Seladin-1 expression in breast cancer cells. 3) Perform functional assays on tumor-specific Seladin-1 mutants.

We amplified the Seladin-1 exons from 60 breast tumor genomic DNA samples (all were obtained from a commercial source (Oncomatrix, Inc.)), followed by the sequence analysis of the open reading frames. As a result, we did not encounter mutations in the coding region and exon-intron junctions (other than known single nucleotide polymorphisms (SNPs)). A single new high frequency insertion/deletion polymorphism was found in the non-coding region. We analyzed the matching DNA from unaffected organs of the same patients. In cases showing this particular SNP, both tumor material and patients DNA obtained from unaffected tissues had the same polymorphism, suggesting that it is not associated with breast cancer. We isolated RNAs from human breast cell lines MCF7, BTK20, T47D, HBL100, MDA-MB-231, MDA-MB-468 and MCF10A. Total RNA isolation was done with Trizol (Invitrogen) as described by the manufacturer. Seladin-1 probe labeling and hybridization was done using commercial kits. As a control, we used RNA from three different batches of normal mammary epithelial cells (HMEC).

When RNAs from 10 primary breast cancer cell lines were analyzed, the conclusions on Seladin-1 gene expression was that this gene is usually expressed at high level in breast cancer cells (with the exception of MDA-MB-231). Interestingly, expression of Seladin-1 in MDA-MB-231 cell line was subjected to regulation by 5aza-2'-deoxycytidine (5AzaDC), a demethylating agent. An increase in Seladin-1 expression in response to 5AzaDC indicates that this putative cell line was a candidate for promoter methylation studies using methylation-specific PCR and bisulfite sequencing.

Since we did not detect breast tumor-specific Seladin-1 mutants (in the coding region), we concentrated on wild type Seladin-1 function and found that Seladin-1 had inhibitory effect on growth of some breast cancer cells (notably MDA-MB-231) expressing it at the low levels, while having no affect on the others: T47D or MDA-MB-468. The work on characterizing the inhibitory function of Seladin-1 in breast tumor cells is currently ongoing.

REFERENCES:

1. Wu, C., Miloslavskaya, I, Demontis, S., Maestro, R., and Galaktionov, K.. *Nature* 2004, **432**, 640-645.
2. Yamashita, H., Nishio, M., Toiyama, T., Sugiura, H., Zhang, Z, Kobayashi, S., and Iwase, H. Coexistence of HER2 over-expression and p53 protein accumulation is a strong prognostic molecular marker in breast cancer. *Breast Cancer Res.* 2004, **6**, R24-R30. Epub 2003 Nov 07.
3. Waterham, H.R. et al. Mutations in the 3beta-hydroxysterol Delta24-reductase gene cause desmosterolosis, an autosomal recessive disorder of cholesterol biosynthesis. *Am. J Hum. Genet.* 2001, **69**, 685-694.